ADVERTISING

GROUP MEMBERS:
ATENA JAFARI PARSA
AVA ARABI
PEYMAN KHODABANDEHLOUEI

INTRODUCTION

The effect of three of the most common (and classic) types of advertisements:

- TV Ads
- Radio Ads
- Newspaper Ads

on companies' sales




DATASET


- The dataset chosen is

 Advertising Dataset of Sales
 with respect to TV, radio and
 newspaper advertisements
 budgets <u>Advertising Dataset</u>
 | <u>Kaggle</u>.
- The sales are in thousands of units and the budget is in thousands of dollars.


a several data points of each variable

Data Analysis







TV ADVERTISEMENTS BUDGET VS SALES



Strong Positive Linear Relationship



Scatter Plot with Linear Regression | Scatter Plot with Linear Regression |




RADIO AND NEWSPAPER **ADVERTISEMENTS BUDGET VS SALES**


No clear relationship

SAMPLE STATISTICS

- A random sample of size 30
- In both the sample and the population: The highest sales' corresponding TV ads budget of \$200K-\$250K budget of TV advertisements
- The lowest sales correspond to TV ads budget larger than \$250K.
- The other bins relatively show a pattern similar to the population histogram as well.



CDF of Standard Normal Distribution



NORMALITY

The Shapiro-Wilk test proved normal distribution for sales population.

Find x as the value of sales for which the z-value of the data points below it (less than x) would make up 0.4 of the data.


$$P(Z \le x) = 0.4$$

HYPOTHESIS TEST 3 CONFIDENCE INTERVALS

- Claim: The population mean of sales is not equal to 15.13. (alpha = 0.05, n = 30)
- H0: mu = 15.13
- Ha: mu != 15.13
- $n = 30 \rightarrow CLT \rightarrow normally distributed$
- qnorm() and pnorm() functions


p-value: 0.8764363

Conclusion: Not enough evidence to reject H0



HYPOTHESIS TEST 2

- Claim: The population mean of Radio is greater than 23. (alpha = 0.05, n = 20)
- H0: mu = 23
- Ha: mu > 23
- t.test() function
- P-value = 0.2074528




DETECTING THE DISTRIBUTION

- Goodness of fit tests
- The Chi-squared test to see whether the TV Budget population is normally distributed. (alpha = 0.05)
- H0: The distribution is normal.
- Ha: The distribution is not normal.
- Chisq.test() function
- P-value < 2.2e-16 < alpha
- Reject H0 in favor of Ha
- TV Budget population is not normally distributed.

LINEAR REGRESSION

- TV: \$230.1K, Radio: \$37.8K, Newspaper: \$69.2K. The predicted sale?
- Fit the model.
- predict()
- Residual error?
- Observed predicted = |22.1 21.2| = 0.9



ANOVA

- Claim: There is a significant difference in the mean values of the three advertising methods. (alpha = 0.05)
- H0: There is no significant difference in the means.
- Ha: There is a significant difference in the means.
- library("dplyr")
- aov() and summary()
- P-value < alpha
- Conclusion: H0 is rejected in favor of Ha.
- There is a significant difference in the mean values of the three advertising methods.

APPLICATIONS OF NONPARAMETRIC TESTS

- Single Sample Sign Test: Is there a significant difference in the sample median from the hypothesized value? (sample sales)
- H0: There is not a significant difference in the sample median from the hypothesized value.
- Ha: There is a significant difference in the sample median from the hypothesized value.
- P-value < alpha
- Conclusion: Reject H0 in favor of Ha.
- There is a significant difference in the sample median from the hypothesized value

THANK YOU FOR LISTENING!

